# **Cardiovascular Topics**

# Right atrial strain in a normal adult African population according to age

Nyange Mushitu, Ruchika Meel

# Abstract

**Background:** Right atrial longitudinal strain (RALS) is a useful parameter to define right atrial (RA) subclinical dysfunction prior to changes in RA dimension and volume. We sought to establish normal values for RALS in a sub-Saharan African population.

**Methods:** This was a retrospective, cross-sectional study from 2017 to 2019 of 100 normal individuals. All echocardiographic measurements were done as per the standard guidelines.

**Results:** Mean RALS was  $32.7 \pm 10.5\%$ . The mean RA volume indexed to body surface area was  $19.5 \pm 5.7$  ml/m<sup>2</sup>. There was a negative correlation between RALS and age but it was not statistically significant (r = -0.15, p = 0.129). Males had a tendency towards higher RA volume indexed and RALS measurements compared to females ( $20.8 \pm 6.3$  and  $18.7 \pm 5.2$  ml/m<sup>2</sup>, p = 0.07;  $34.6 \pm 9.6$  and  $31.4 \pm 10.9\%$ , p = 0.141, respectively). Body mass index was an independent predictor of RALS (r = -0.43, p = 0.003).

**Conclusion:** We have provided normative data for RALS in an African population. This study provides a platform for future larger studies on RALS.

Keywords: right atrium, right atrial longitudinal strain, age, African, echocardiography

| Submitted 25/2/22, accepted 7/2/23        |                |
|---|----------------|
| Cardiovasc J Afr 2023; online publication | www.cvja.co.za |

DOI: 10.5830/CVJA-2023-004

In the last 20 years, there has been great interest in research on right atrial (RA) function, because recent data have demonstrated that assessment of RA volumetric parameters on echocardiography is an essential predictor of morbidity and mortality in various cardiovascular disorders.<sup>13</sup>

#### Department of Internal Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

Nyange Mushitu, MB ChB, nyangemush@gmail.com

Division of Cardiothoracic Surgery, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

Ruchika Meel, MB ChB, MMed, Cert Cardiology (SA), PhD, FEACVI, FESC, RuchikaMeel@gmail.com

There are several imaging studies that have described RA anatomy and function. RA dimensions, volume and strain have been studied using the gold standard, cardiac magnetic resonance (CMR), computed tomography, echocardiography (two- and three-dimensional) and recently speckle-tracking echocardiography (STE). Two-dimensional echocardiography (2DE) has become the most clinically relevant non-invasive technique for evaluating the right atrium.<sup>46</sup>

It is important to define normative values of RA strain using 2DE, because RA subclinical dysfunction, as measured by strain, has been observed in several cardiovascular disorders prior to changes in the traditional indices of RA and right ventricular (RV) function such as volume, size and ejection fraction. Such cardiovascular disorders include pulmonary arterial hypertension (PAH), coronary artery disease (CAD) and heart failure with reduced ejection fraction (HFrEF). RA strain had additive prognostic value to other clinical measures, including RV strain, RA area and RA pressure in patients with PAH.<sup>7-11</sup> The difference between normal and abnormal RA dimension and function is therefore clinically pertinent.

Most of the available studies on the parameters of RA dimension in a normal population depend on data from North America and Europe and are in line with the guidelines of the American Society of Echocardiography (ASE) and European Society of Cardiology (ESC).<sup>12-15</sup> These values do not effectively represent the diverse racial and ethnic groups of the world.

The study by Soulat-Dufour *et al.* with the World Alliance Societies of Echocardiography (WASE) have suggested that there might be significant differences in normal values among different populations.<sup>16</sup> Currently, limited data exist regarding RA volume (RAV), size and strain in a normal black African adult population.

Age-related changes in vascular and cardiac function contribute to cardiovascular mortality. Aging is associated with abnormalities in left-sided functional parameters. However, studies on age-related changes in right-sided functional parameters are scarce.<sup>17-19</sup> Therefore, in this study, we sought to establish normal values for RAV and RA longitudinal strain (RALS), and its correlation with age, in a sub-Saharan black African population, using 2DE and STE.

#### Methods

This study was a retrospective analysis of echocardiographic findings in healthy normal controls that formed part of a study (M170389) conducted at Chris Hani Baragwanath Academic Hospital (CHBAH). It is a secondary analysis of the data collected in these healthy controls.



Fig. 1. Apical four-chamber view in a normal participant, showing measurement of RA volume using the discs method.

The study was approved by the University of the Witwatersrand ethics committee and conforms to the principles outlined in the Declaration of Helsinki. Permission to use data from the parent study was obtained. The study was approved by the Human Research Ethics Committee of the University of the Witwatersrand (clearance certificate number: M 200822).

Echocardiograms were obtained according to a standardised protocol on a Phillips iE 33 ultrasound system (Amsterdam, The Netherlands) equipped with an S5-1 transducer that transmits a frequency of 1.7 MHz and receives a frequency of 3.4 MHz. All data were analysed offline after being transferred to an Xcelera workstation (Philips). The RALS analysis was performed separately offline using Philips Q lab 11.0 speckle-tracking software.

Two-dimensional echocardiography was used for measurements of RA dimensions. They were performed in agreement with ASE chamber quantification guidelines of 2015 and the 2010 ASE guidelines on right heart assessment. The RA parameters (area and volume) were measured at the end of systole on an apical four-chamber view modified to optimise the RA visualisation. The RAV was measured using the single-plane method of discs by tracing an outline of the RA blood–tissue interface, ensuring that the RA appendage, superior vena cava and inferior vena cava were excluded. The tricuspid tenting area was also excluded (Fig. 1).<sup>12,14,15</sup>

The RALS was measured by an experienced cardiologist and clinical technologist using STE. For STE using 2D grey-scale echocardiography, apical four-chamber views were captured during breath holding at the end of the expiratory phase for a few seconds, and with an electrocardiogram recording attached. A suitable image of myocardial tissue was obtained completely separated from surrounding structures. Three successive cardiac cycles were recorded and averaged. The frame rate was set between 60 and 80 frames per second. Analysis of specklebased strain was done using Philips Q lab 11.0 speckle-tracking software.

In four-chamber RA focused views, the endocardial surface of the RA was traced manually by a three-point-and-click approach. The system automatically generates an epicardial surface tracing. The region of interest (ROI) was therefore created, composed of seven segments. The ROI was manually adjusted as needed to allow for enough speckle tracking. The software generates the longitudinal curves for each segment with its mean value (Fig. 2).<sup>20-22</sup>

The number of healthy controls who were enrolled in the parent study was 100, after excluding 23 patients who did not meet the study inclusion criteria. Using the power command in Stata, we conducted a one-sample correlation test to estimate the minimum sample required to detect a correlation of at least r = 0.3 between RALS and age. The minimum sample size required was 100. All participants were subdivided into four age groups (group 1: 18–29 years, group 2: 30–39 years, group 3: 40–49 years and group 4:  $\geq$  50 years).

This study is a secondary data analysis, and we utilised data from an existing database. Only variables in our data collection sheet were used in this study.

#### Statistical analysis

Continuous variables are described using the mean and standard deviation, or median and interquartile range when variables were not normally distributed. The independent *t*-test was used to compare means of continuous variables by gender while the Mann–Whitney test was used for a comparison of median values by gender. One-way analysis of variance (ANOVA) was used to



Fig. 2. STE showing decreased peak RA longitudinal strain of 25.5% in an older subject (A), compared to a younger subject of 46.5% (B).

compare means of normally distributed continuous variables by age group or body mass index (BMI) category, while the Kruskal–Wallis test was used for comparison of medians when variables were not normally distributed. Pearson's correlation coefficients determined the association between age and RA parameters using a statistical significance threshold of 0.05.

All statistical analyses were conducted in Stata version 15. Univariate and multivariable linear regression was used to explore the association between RALS and independent variables at presentation [age, gender, BMI, body surface area (BSA) and RV parameters]. Independent variables were selected *a priori* for multivariate analyses. All clinically relevant and/or statistically significant variables on univariate analysis with a *p*-value  $\leq 0.1$  were included in the multivariate analysis.

## **Results**

Baseline characteristics of the overall population of the study are presented in Table 1. A total of 100 participants were included, with a median age of 37.5 years [interquartile range (IQR) 26-46] with 60% female, and 24% of participants were above 50 years of age. Weight differed by age group (p = 0.049) and the median BMI was 28.0 (24.0-34.9) kg/m<sup>2</sup>. A greater percentage of female participants were overweight and obese (50%). There was an age-related increase in BMI of all participants, but it was not statistically significant (p = 0.089).

Two-dimensional echocardiography parameters according to age groups are depicted in Table 2. There were no statistically significant differences with regard to age-related RA volumetric measurements (p = 0.271). RALS had a trend towards decreasing with age but did not achieve statistical significance (p = 0.362). RA volume indexed (RAVI) and RALS had a negative correlation with age but did not reach statistical significance (r = -0.060, p = 0.526; r = -0.153, p = 0.129, respectively) (Fig. 3).

Regarding traditional RV functional parameters such as tricuspid annular plane systolic excursion (TAPSE), RV E', RV A', and RV S') there were no statistically significant differences. RV diastolic function declined with age, with RV E'/A' showing a decrease with age (p = 0.002). Left ventricular (LV) diastolic function parameters showed a decrease with age but remained within ASE guideline-specified normal limits (p < 0.001).

Echocardiographic parameters according to gender are depicted in Table 3. Males had a tendency towards higher RAVI and RALS measurements compared to females  $(20.8 \pm 6.3 \text{ and})$ 

18.7  $\pm$  5.2 ml/m<sup>2</sup>, p = 0.07; 34.6  $\pm$  9.6 and 31.4  $\pm$  10.9%, p = 0.141, respectively). There were no statistically significant differences in RV measurements and functional parameters between males and females. All participants had LV diastolic pulsed-wave (transmitral E and A waves) and tissue Doppler measurements within the normal accepted guideline range except for A wave and A' lateral, which were higher in females compared to males (p = 0.001 and p = 0.004, respectively). Male participants had higher LV end-diastolic and end-systolic dimensions and volumes, as well as LA volume indexed to BSA.

Clinical and echocardiographic indices according to BMI are depicted in Table 4. Twenty-three per cent of participants were overweight and 44% were obese. Obese participants had higher heart rates than non-obese participants (p = 0.002) There were no statistically significant differences with regard to RA volumetric measurements, but RALS had a tendency to decrease with increasing BMI ( $35.7 \pm 9.3\%$ ,  $< 25 \text{ kg/m}^2$ ;  $34.6 \pm 11.4\%$ ,  $< 30 \text{ kg/m}^2$ ; and  $29.9 \pm 10.1\%$ ,  $> 30 \text{ kg/m}^2$ , respectively, p = 0.571). RV E'/A' ratio decreased with increasing BMI (p = 0.020), suggestive of worsening RV diastolic dysfunction with increasing BMI. Left atrial (LA) volume and size increased with an increasing BMI (p = 0.010 and p < 0.001, respectively). Similar to RV diastolic parameters, there was worsening of LV diastolic function with an increase in BMI. Determinants of RA longitudinal strain are depicted in Table 5.

BMI was an independent predictor of RALS (BMI was a statistically significant variable on both univariate and multivariate analyses). BMI negatively correlated with RALS (r = -0.43, p = 0.003). For example, in the adjusted analyses, a unit increase in BMI resulted in a 0.72 decline in RALS. The intraobserver coefficient of variation for peak RALS was 4.6% and the inter-observer variability was 9%.

### Discussion

This is the first study to provide normative age-related data for RALS in a sub-Saharan African population. Additionally, we have also provided complementary data regarding RAVI according to age and gender in this population. RALS had a tendency to decrease with age concurrent with a decline in RV diastolic function despite no alteration in RAVI, and BMI was an important independent predictor of RALS. Even though there was a tendency of RAVI to be higher in males, RAVI and RALS were not influenced significantly by gender.

| Clinical parameters                  | Total (20-62 years) (n = 100) | Group 1<br>(18–29 years)<br>(n = 27) | Group 2 $(30-39 years)$ $(n = 28)$ | Group 3<br>(40–49 years)<br>(n = 21) | Group 4<br>( $\geq 50$ years)<br>(n = 24) | p-value*<br>(ANOVA)# |
|--------------------------------------|-------------------------------|--------------------------------------|------------------------------------|--------------------------------------|---|----------------------|
| Age (years)                          | 37.5 (29.0-48.0)              | 25.0 (23.0-28.0)                     | 34.0 (33.0-37.0)                   | 45.0 (42.0-47.0)                     | 54.0 (51.0-57.5)                          | < 0.001              |
| Height (cm)                          | $160.7 \pm 7.3$               | 161.6 ± 8.2                          | $160.6 \pm 6.5$                    | $162.0 \pm 6.4$                      | 158.8 ± 7.9                               | 0.499                |
| Weight (kg)                          | 73.0 (65.0-85.0)              | 67.0 (59.0-75.0)                     | 73.0 (65.0-86.5)                   | 80.0 (64.0-87.0)                     | 78.3 (69.3-84.5)                          | 0.049                |
| Female, $n$ (%)                      | 60 (60.0)                     | 13 (48.2)                            | 18 (64.3)                          | 10 (47.6)                            | 19 (79.2)                                 | 0.078                |
| Body mass index (kg/m <sup>2</sup> ) | 28.0 (24.0-34.9)              | 24.0 (22.5–29.3)                     | 29.1 (24.2-35.4)                   | 29.0 (26.6-34.3)                     | 30.8 (27.1-36.6)                          | 0.089                |
| Body surface area (m <sup>2</sup> )  | $1.8 \pm 0.2$                 | $1.7 \pm 0.2$                        | $1.8 \pm 0.2$                      | $1.8 \pm 0.2$                        | $1.8 \pm 0.1$                             | 0.168                |
| Systolic blood pressure (mmHg)       | $126.8 \pm 12.4$              | $123 \pm 11.6$                       | $126 \pm 14.0$                     | $127 \pm 9.1$                        | $130.8 \pm 13.2$                          | 0.205                |
| Diastolic blood pressure (mmHg)      | $79.3 \pm 10.3$               | $76.9 \pm 10.9$                      | $79.2 \pm 11.6$                    | $80.5 \pm 9.3$                       | $80.9 \pm 9.0$                            | 0.498                |
| Heart rate (bpm)                     | 72.0 (62.5-82.0)              | 67.0 (57.0-81.0)                     | 75.5 (67.0-85.5)                   | 70.0 (61.0-82.0)                     | 71.0 (62.5-81.5)                          | 0.217                |

Kruskall–Wallis p-value for non-normally distributed variables. \*Statistical significance denoted by p < 0.05.

|   | Table 2. Echocardiogra    | aphic parameters ac       | cording to the age o                            | f the study group        |                                   |           |
|---|---------------------------|---------------------------|---|--------------------------|-----------------------------------|-----------|
|   | Total<br>(20–62 years)    | Group 1<br>(18–29 years)  | Group 2<br>(30–39 years)                        | Group 3<br>(40–49 years) | Group 4 $(\geq 50 \text{ years})$ | p-value*  |
| Variable  | (n = 100)                 | (n = 27)                  | (n = 28)  | (n = 21)                 | (n = 24)                          | (ANOVA)#  |
| RA parameters   |                           |                           |   |                          |                                   |           |
| RA volume (ml)  | $34.8 \pm 10.7$           | $35.3 \pm 11.7$           | $33.7 \pm 8.1$                                  | $36.1 \pm 13.8$          | $34.3 \pm 9.0$                    | 0.865     |
| RA volume index (ml/m <sup>2</sup> )  | $19.5 \pm 5.7$            | $20.2 \pm 5.8$            | $19.0 \pm 4.9$                                  | $19.8 \pm 7.2$           | $19.1 \pm 5.3$                    | 0.271     |
| RA length (mm)  | $44.5 \pm 5.2$            | $44.5 \pm 5.1$            | $44.6 \pm 3.9$                                  | $45.3 \pm 6.0$           | $43.7 \pm 6.0$                    | 0.679     |
| RA width (mm)   | $36.3 \pm 5.6$            | $36.7 \pm 5.9$            | $36.9 \pm 5.6$                                  | $36.4 \pm 6.3$           | $34.9 \pm 5.3$                    | 0.588     |
| RALS (%)  | $32.7 \pm 10.5$           | $34.7 \pm 10.8$           | $32.8 \pm 11.7$                                 | $33.5 \pm 7.3$           | $29.6 \pm 11.1$                   | 0.362     |
| RV functional parameters  |                           |                           |   |                          |                                   |           |
| TAPSE (mm)  | 19.0 (17.0-22.0)          | 17.0 (16.0–19.0)          | 19.0 (17.0-21.5)                                | 19.0 (18.0-22.0)         | 19.5 (17.0-25.5)                  | 0.099     |
| PASP (mmHg)   | 15.0 (13.5-20.0)          | 15.0 (13.5-20.0)          | 17.0 (15.0-20.0)                                | 18.0 (13.0-26.0)         | 15.0 (10.0-18.0)                  | 0.465     |
| RV base (mm)  | 31.0 (27.5-33.2)          | 32.0 (28.0-34.0)          | 30.0 (27.1-32.4)                                | 31.4 (28.5–32.9)         | 30.0 (26.6-33.9)                  | 0.455     |
| RV S' (cm/s)  | $11.1 \pm 2.1$            | $10.7 \pm 1.9$            | $11.3 \pm 2.4$                                  | $11.4 \pm 2.0$           | $11.0 \pm 2.0$                    | 0.677     |
| RV E' (cm/s)  | 11.0 (8.9–13.0)           | 12.9 (9.5–15.9)           | 11.0 (9.7–12.6)                                 | 11.1 (8.0-12.5)          | 10.0 (7.6–11.5)                   | 0.085     |
| RV A' (cm/s)  | 9.7 (8.4-12.1)            | 8.5 (7.9-10.6)            | 9.9 (8.7-11.6)                                  | 11.0 (9.0-13.4)          | 10.7 (9.0-12.8)                   | 0.015     |
| RV E'/A'  | 1.1 (0.8–1.4)             | 1.4 (1.1–1.9)             | 1.0 (0.9–1.3)                                   | 0.9 (0.8–1.3)            | 0.8 (0.7-1.1)                     | 0.002     |
| LV measurements   |                           |                           |   |                          |                                   |           |
| LV systolic diameter (mm)   | 29.0 (26.0-31.5)          | 29.0 (26.0-32.0)          | 28.5 (26.0-30.0)                                | 30.0 (28.0-32.0)         | 27.0 (23.5-31.0)                  | 0.213     |
| LV diastolic diameter (mm)  | $42.3 \pm 5.1$            | $42.8 \pm 6.0$            | $41.9 \pm 4.2$                                  | $44.0 \pm 4.2$           | $40.5 \pm 5.3$                    | 0.124     |
| End-diastolic volume index (ml/m <sup>2</sup> )                                     | $86.2 \pm 23.1$           | $89.3 \pm 26.6$           | $89.5 \pm 24.2$                                 | $91.0 \pm 16.9$          | $74.8 \pm 19.7$                   | 0.048     |
| End-systolic volume index (ml/m <sup>2</sup> )                                      | 18.2 (14.5-22.6)          | 20.1 (18.3-26.0)          | 18.3 (16.3-25.7)                                | 17.4 (13.2–19.3)         | 14.3 (11.1–17.2)                  | < 0.001   |
| LA volume (ml)  | 36.8 (27.7-46.8)          | 41.0 (23.0-50.0)          | 35.0 (26.0-45.9)                                | 41.4 (29.0-44.4)         | 37.0 (31.5-48.0)                  | 0.630     |
| LA volume index (ml/m <sup>2</sup> )  | 21.5 (15.2-26.4)          | 23.8 (14.8-28.5)          | 19.6 (13.4-23.8)                                | 22.7 (14.5-25.7)         | 20.0 (17.3–27.8)                  | 0.574     |
| LA size (mm)  | 32.0 (28.0-35.3)          | 30.8 (27.0-33.4)          | 32.0 (29.0-35.8)                                | 33.0 (27.8–38.0)         | 33.0 (29.5–38.0)                  | 0.068     |
| Ejection fraction (%)   | $62.5 \pm 7.1$            | $60.5 \pm 7.3$            | $61.7 \pm 6.9$                                  | $65.5 \pm 6.2$           | $63.0 \pm 7.5$                    | 0.104     |
| Posterior wall diameter (mm)  | 8.0 (7.0-9.5)             | 7.0 (7.0-8.0)             | 8.0 (7.0-9.0)                                   | 9.0 (8.0-10.0)           | 9.0 (8.0-11.0)                    | 0.001     |
| LV diastolic parameters   |                           |                           |   |                          |                                   |           |
| E wave (cm/s)   | $80.6 \pm 19.7$           | 89.1 ± 18.1               | $82.1 \pm 17.6$                                 | $76.5 \pm 20.9$          | $72.7 \pm 19.7$                   | 0.017     |
| A wave (cm/s)   | $56.2 \pm 15.1$           | 49.1 ± 15.3               | $57.3 \pm 13.8$                                 | $54.2 \pm 13.1$          | $64.9 \pm 14.4$                   | 0.002     |
| E/A ratio   | 1.4 (1.1–1.8)             | 1.8 (1.6-2.2)             | 1.4 (1.2–1.8)                                   | 1.4 (1.2–1.5)            | 1.1 (1.0-1.3)                     | < 0.001   |
| E' medial (cm/s)  | 9.5 (7.5–11.8)            | 11.7 (10.1–13.6)          | 9.1 (7.7–11.8)                                  | 9.0 (7.0–10.0)           | 7.7 (5.8–9.0)                     | < 0.001   |
| E' lateral (cm/s)   | $13.4 \pm 3.3$            | $16.0 \pm 3.1$            | $13.9 \pm 2.9$                                  | $12.0 \pm 2.4$           | $11.1 \pm 2.6$                    | < 0.001   |
| A' lateral (cm/s)   | $8.7 \pm 2.7$             | $7.0 \pm 1.9$             | $9.0 \pm 2.4$                                   | $9.6 \pm 2.6$            | $9.6 \pm 2.9$                     | < 0.001   |
| E/E' lateral ratio  | 5.9 (5.0-7.0)             | 5.8 (4.2-7.0)             | 5.8 (5.0-6.4)                                   | 5.9 (5.0-7.0)            | 6.2 (5.0-8.2)                     | 0.575     |
| Data reported as means $\pm$ SD or media<br>atrium longitudinal strain: RV right ve | an (IQR). IQR, interquart | ile range; LA, left atria | ıl; PASP, pulmonary a<br>d annular plane systol | rtery systolic pressure; | RA, right atrial; RAI             | .S. right |

"Independent t-test, p-value of (Mann–Whitney test, p-value for non-normally distributed variables). \*Statistical significance denoted by p-values < 0.05

Previous studies using 2DE by Wang *et al.* in 1984, and Kou *et al.* from the Normal Reference Ranges for Echocardiography (NORRE) study in 2013, were focused on Caucasians populations,

and were in line with the ASE and ESC 2015 chamber quantification guideline document, which states that RAVI should be  $< 33 \text{ ml/m}^2$  (or  $< 35 \text{ ml/m}^2$  in men and  $< 31 \text{ ml/m}^2$  in women).<sup>23,24</sup>



r = -0.153, p = 0.129, respectively).

| of   | study participa   | nts according to  | o gender   |                           |
|--|---|---|--|---------------------------|
|  |   |   |  | p-value                   |
| Variables  | Total ( $n = 100$ )   | Males (n = 40)  | Females $(n = 60)$   | (ANOVA                    |
| Clinical parameters  |   |   |  |                           |
| Age (years)  | 37.5 (29.0–48.0)  | 34.0 (27.0–45.5)  | 39.0 (32.5–51.0)   | 0.055                     |
| Weight (kg)  | 73.0 (65.0–85.0)  | 68.0 (63.3–79.5)  | 79.0 (66.0–94.0)   | 0.016                     |
| Body mass index (kg/m <sup>2</sup> )   | 28.0 (24.0–34.9)  | 24.1 (22.5–29.6)  | 31.2 (26.6–37.8)   | < 0.001                   |
| Body surface area (m <sup>2</sup> )  | $1.8 \pm 0.2$   | $1.8 \pm 0.1$   | $1.8 \pm 0.2$  | 0.783                     |
| Systolic blood<br>pressure (mmHg)  | 126.8 ± 12.4  | $127.2 \pm 12.2$  | 126.3 ± 12.5   | 0.737                     |
| Diastolic blood<br>pressure (mmHg)   | 79.3 ± 10.3   | 78.5 ± 11.8   | 79.8 ± 9.3   | 0.535                     |
| Heart rate (bpm)   | 72.0 (62.5-82.0)  | 65.0 (59.0–76.5)  | 76.0 (67.0-82.5)   | 0.002                     |
| RA parameters  |   |   |  |                           |
| RA volume (ml)   | $34.8 \pm 10.7$   | $37.0 \pm 11.8$   | $33.3 \pm 9.5$   | 0.090                     |
| RA volume index<br>(ml/m <sup>2</sup> )  | $19.5 \pm 5.7$  | 20.8 ± 6.3  | $18.7 \pm 5.2$   | 0.070                     |
| RA width (mm)  | $36.3 \pm 5.6$  | $37.9 \pm 6.0$  | $35.2 \pm 5.2$   | 0.018                     |
| RA length (mm)   | $44.5 \pm 5.2$  | $44.2 \pm 5.6$  | $44.7 \pm 4.9$   | 0.679                     |
| RALS (%)   | $32.7\pm10.5$   | $34.6 \pm 9.6$  | $31.4\pm10.9$  | 0.141                     |
| RV functional paramet  | ers   |   |  |                           |
| TAPSE (mm)   | 19.0 (17.0–22.0)  | 19.0 (17.0-22.0)  | 18.5 (17.0–21.5)   | 0.659                     |
| PASP (mmHg)  | 15.0 (13.5–20.0)  | 15.0 (12.0–17.0)  | 17.0 (15.0–20.0)   | 0.056                     |
| RV E' (cm/s)   | 11.0 (8.9–13.0)   | 11.2 (9.0–12.9)   | 10.9 (8.6–13.2)  | 0.699                     |
| RV A' (cm/s)   | 9.7 (8.4–12.1)  | 9.1 (8.4–11.7)  | 10.2 (8.7–12.8)  | 0.155                     |
| RV S' (cm/s)   | $11.1 \pm 2.1$  | $10.9 \pm 2.2$  | $11.2 \pm 1.9$   | 0.406                     |
| RV base (mm)   | 31.0 (27.5–33.2)  | 31.6 (27.8–33.2)  | 30.3 (27.1–33.3)   | 0.499                     |
| RV E/A'  | 1.1 (0.8–1.4)   | 1.0 (0.8–1.3)   | 1.1 (0.8–1.5)  |                           |
| LV measurements  |   |   |  |                           |
| LV systolic diameter (mm)  | 29.0 (26.0–31.5)  | 30.0 (27.5–32.0)  | 27.0 (25.5–30.0)   | 0.006                     |
| LV diastolic diameter (mm)   | 42.3 ± 5.1  | 44.2 ± 5.7  | 41.0 ± 4.2   | 0.001                     |
| End-diastolic volume index (ml/m <sup>2</sup> )  | $86.2 \pm 23.1$   | 94.5 ± 26.2   | 80.8 ± 19.2  | 0.003                     |
| End-systolic volume index (ml/m <sup>2</sup> )   | 18.2 (14.5–22.6)  | 16.9 (14.2–19.4)  | 19.8 (17.2–24.7)   | 0.012                     |
| LA volume index (ml/m <sup>2</sup> )   | 21.5 (15.2–26.4)  | 22.6 (16.1–28.0)  | 19.1 (13.3–24.6)   | 0.060                     |
| LA size (mm)   | 32.0 (28.0–35.3)  | 32.0 (28.3–34.0)  | 32.2 (27.8–36.6)   | 0.441                     |
| Ejection fraction (%)  | $62.5 \pm 7.1$  | $62.2 \pm 7.8$  | $62.7\pm6.7$   | 0.753                     |
| Posterior wall<br>diameter (mm)  | 8.0 (7.0–9.5)   | 8.0 (7.0–9.5)   | 8.0 (7.0–9.5)  | 0.516                     |
| LV diastolic parameter   | s   |   |  |                           |
| E wave (cm/s)  | $80.6 \pm 19.7$   | $78.2 \pm 16.1$   | $82.2\pm21.8$  | 0.320                     |
| A wave (cm/s)  | $56.2 \pm 15.1$   | $50.3 \pm 15.6$   | $60.2 \pm 13.5$  | 0.001                     |
| E/A ratio  | 1.4 (1.1–1.8)   | 1.5 (1.2–1.9)   | 1.4 (1.1–1.4)  | 0.075                     |
| E' medial (cm/s)   | 9.5 (7.5–11.8)  | 9.7 (7.9–12.0)  | 9.4 (7.1–11.5)   | 0.429                     |
| E' lateral (cm/s)  | $13.4 \pm 3.3$  | $13.8 \pm 3.0$  | $13.1 \pm 3.5$   | 0.287                     |
| E/E' lateral ratio   | 5.9 (5.0-7.0)   | 5.7 (5.0-6.8)   | 6.0 (5.0-7.1)  | 0.215                     |
| Data reported as mean<br>IQR, interquartile rang<br>right atrial; RALS, righ<br>deviation; TAPSE, trict<br>"Independent <i>t</i> -test, <i>p</i> -v<br>distributed variables. *{ | s $\pm$ SD or median<br>ge; LA, left atrial;<br>nt atrial longitudin<br>uspid annular pla<br>value or Mann–W<br>Statistical signific: | (IQR).<br>PASP, pulmonary<br>nal strain; RV, rig<br>ne systolic excurs:<br>'hitney test, <i>p</i> -valu<br>ance denoted by <i>t</i> | y artery systolic pr<br>ht ventricle; SD, st<br>ion.<br>ie for non-normal<br>y-values < 0.05 | essure; R<br>andard<br>ly |

In our study, we found that while the lower limits were similar to those of ASE and ESC guidelines, the upper margins were much lower (20.8  $\pm$  6.3 ml/m<sup>2</sup> in male and 18.7  $\pm$  5.2 ml/m<sup>2</sup> in females). Studies in African populations by Nel *et al.* and in Asian subjects by Karki *et al.* observed the same pattern.<sup>25,26</sup> Soulat-Dufour *et al.*, in the WASE study with 2 008 healthy adult individuals around the world, has shown that generally, Asian

subjects had lower BSA compared to subjects in non-Asian countries.<sup>16</sup>

Ethnicity may cause variations in genetics and parameters of BSA and BMI and may directly influence echocardiographic measures. Therefore, correlation between ethnicity and RA volumetric parameters is important.

In this study, there was a trend towards higher RAVI in males compared to females but no influence of age on RAVI was noted. The aforementioned finding was akin to the recent study by Nel *et al.*<sup>25</sup> and the WASE study pertaining to RAVI in a normal population.<sup>16</sup> Grünig *et al.* and D'Ascenzi *et al.* have shown that males had a larger RA area compared to females, explaining the volume differences between genders.<sup>27,28</sup> D'Oronzio *et al.*, Wang *et al.* and Peluso *et al.*, using 2DE and 3DE, respectively, did not find any correlation between atrial volume and ageing, for reasons that are not fully understood.<sup>23,29,30</sup>

In the current study, the mean RALS was  $32.7 \pm 10.5\%$ , which was lower than the value reported by Padeletti *et al.* and D'Ascenzi *et al.*<sup>22,28</sup> In this Italian study comprising 84 and 74 subjects, RALS was documented using the 2D STE, and the mean RALS was  $49 \pm 13$  and  $48 \pm 12.68\%$ , respectively.<sup>22,28</sup> These studies were done in a group of subjects with an overall lower BMI ( $22.4 \pm 3.5 \text{ kg/m}^2$ ), in contrast to our study ( $24-34.9 \text{ kg/m}^2$ ), and this is likely the explanation for higher RALS in their cohort.

Obesity is a health problem of growing significance all over the world; its prevalence is increasing in both developed and developing countries. According to World Health Organisation data, 39% of the global population above 18 years of age are overweight and of these, 13% are obese. In Africa there is a significant trend towards obesity, with increments in BMI documented in both genders.<sup>31</sup> Micklesfield *et al.* studied a South Africa population from Soweto and demonstrated a significant gender difference with regard to BMI, with women being markedly more overweight and obese.<sup>32</sup>

In our study, 67% of normal volunteers had a BMI above 25 kg/m<sup>2</sup>, which occurred more commonly in women (49%). BMI was the only independent predictor of RALS in this study. An inverse relationship was noted between BMI and RALS in the current study. A recent study by Chirinos *et al.* quantified LA strain and strain rate using STE in 1 531 middle-aged community-based participants enrolled in the Asklepios study. They demonstrated that longitudinal LA strain measured using STE decreased with elevated BMI.<sup>33</sup>

Obesity has been shown to have many effects on cardiovascular structure and function. Excess adiposity imposes an increased metabolic demand on the body and cardiac output. Total blood volumes are elevated in obesity, leading to a hyperdynamic circulation, which causes LV and RV structural changes, with the resultant increased ventricular mass and cavity dilatation. Obesity is the main cause of tissues fibrosis.<sup>34</sup>

Sokmen *et al.* and Csige *et al.* showed that uncomplicated obesity was associated with RV and RA dilatation, and increased thickness of the RV free wall. Also, these structural indices were found to be positively correlated with BMI.

Myocardial fat accumulation as a consequence of obesity may cause atrial interstitial fibrosis and subsequently atrial dilatation and stiffness.<sup>35,36</sup> This may be a possible mechanism explaining the inverse relationship between BMI and RALS in this study. Obesity may result in RA fibrosis, stiffness with a consequent

|   | Table 4. Clinical and echoc   | ardiographic indices a   | ccording to body mass ind                               | ex                                    |                  |
|---|---|--|---|---------------------------------------|------------------|
| V   | $BMI kg/m^2$  | $BMI < 25 \ kg/m^2$  | $BMI 25 \ to < 30 \ kg/m^2$                             | $BMI \ge 30 \ kg/m^2$                 | p-value*         |
| Clinical parameters   | (11 - 100)  | (11 - 32)  | (11 - 23)   | (11 - 44)                             | (ANOVA)          |
|   | 27.5 (20.0, 48.0)   | 20.0 (24.5, 28.5)  | 40.0 (20.0, 48.0)                                       | 44.0 (24.0, 51.0)                     | < 0.001          |
| Age (years)   | 57.5(29.0-40.0)<br>72.0(65.0.85.0)  | 50.0(24.3-50.5)  | 40.0 (30.0-40.0)  | 44.0 (54.0-51.0)<br>86.5 (82.0, 00.5) | < 0.001          |
| E-male in (9/)  | (0.0-03.0)  | 10(21.2)   | 15(65.2)  | 80.3 (82.0-99.3)<br>24 (77.2)         | < 0.001<br>0.078 |
| Female, $n(\gamma_0)$   | 00(00.0)  | 10(51.3)<br>$1.7 \pm 0.1$  | 13(03.2)<br>$1.7 \pm 0.1$                               | 34(7/.3)                              | 0.078            |
| Body surface area (m <sup>-</sup> )   | $1.8 \pm 0.2$   | $1./ \pm 0.1$  | $1.7 \pm 0.1$   | $1.9 \pm 0.2$                         | 0.009            |
| Systolic blood pressure (mmHg)  | $120.8 \pm 12.4$  | $123.0 \pm 12.9$   | $128.1 \pm 10.7$  | $128.5 \pm 12.8$                      | 0.598            |
| Diastolic blood pressure (mmHg)   | $79.3 \pm 10.3$   | $76.6 \pm 10.8$  | $79.3 \pm 10.8$   | 81.2 ± 9.7                            | 0.760            |
| Heart rate (bpm)  | 72.0 (62.5-82.0)  | 65.5 (58.0–77.0)   | 65.0 (57.0-81.0)  | 77.0 (70.5-82.5)                      | 0.002            |
| RA parameters   |   |  |   |                                       |                  |
| RA volume (ml)  | $34.8 \pm 10.7$   | $35.1 \pm 11.6$  | $30.7 \pm 9.2$  | $36.7 \pm 10.2$                       | 0.513            |
| RA volume index (ml/m <sup>2</sup> )  | $19.5 \pm 5.7$  | $20.8 \pm 5.9$   | $18.0 \pm 5.9$  | $19.4 \pm 5.4$                        | 0.822            |
| RA length (mm)  | $44.5 \pm 5.2$  | $42.9 \pm 5.7$   | $43.4 \pm 4.0$  | $46.2 \pm 5.0$                        | 0.226            |
| RA width (mm)   | $36.3 \pm 5.6$  | $37.2 \pm 5.5$   | $33.2 \pm 5.1$  | $37.3 \pm 5.8$                        | 0.817            |
| RALS (%)  | $32.7 \pm 10.5$   | $35.7 \pm 9.3$   | $34.6 \pm 11.4$   | $29.9 \pm 10.1$                       | 0.571            |
| RV functional parameters  |   |  |   |                                       | 1                |
| TAPSE (mm)  | 19.0 (17.0-22.0)  | 19.0 (17.0–22.5)   | 19.0 (16.0-23.2)  | 18.5 (17.0–21.5)                      | 0.898            |
| PASP (mmHg)   | 15.0 (13.5–20.0)  | 15.0 (15.0–25.0)   | 17.5 (15.0-20.0)  | 15.0 (12.0–18.0)                      | 0.091            |
| RV E' (cm/s)  | 11.0 (8.9–13.0)   | 12.0 (10.1–13.6)   | 11.7 (8.9–13.6)   | 10.2 (8.0–11.6)                       | 0.114            |
| RV A' (cm/s)  | 9.7 (8.4–12.1)  | 9.0 (7.6–11.8)   | 10.2 (8.7–11.3)   | 10.3 (8.8–13.0)                       | 0.081            |
| RV E'/ A'   | 1.1 (0.8–1.4)   | 1.2 (0.9–1.6)  | 1.0 (0.8–1.5)   | 0.9 (0.7-1.2)                         | 0.020            |
| LV measurements   |   |  |   |                                       |                  |
| LV systolic diameter (mm)   | 29.0 (26.0-31.5)  | 29.5 (25.5-32.0)   | 29.0 (26.0-30.0)  | 29.0 (26.0-32.0)                      | 0.730            |
| LV diastolic diameter (mm)  | $42.3 \pm 5.1$  | $42.1 \pm 5.2$   | $42.5 \pm 4.7$  | $42.4 \pm 5.3$                        | 0.824            |
| End-diastolic volume index (ml/m2)  | $86.2 \pm 23.1$   | 88.0 ± 23.3  | 88.6 ± 24.6   | $84.0 \pm 22.7$                       | 0.913            |
| End-systolic volume index (ml/m <sup>2</sup> )  | 18.2 (14.5-22.6)  | 20.0 (16.1-23.5)   | 18.1 (15.3–19.9)  | 17.1 (13.4–21.1)                      | 0.220            |
| LA volume index (ml/m <sup>2</sup> )  | 21.5 (15.2-26.4)  | 20.1 (13.0-23.8)   | 22.6 (16.7-25.8)  | 22.8 (16.2-30.5)                      | 0.242            |
| LA size (mm)  | 32.0 (28.0-35.3)  | 28.0 (25.0-32.0)   | 30.3 (27.0-32.0)  | 35.8 (33.0-38.5)                      | < 0.001          |
| Eiection fraction (%)   | $62.5 \pm 7.1$  | $62.7 \pm 5.8$   | $64.3 \pm 7.9$  | $61.4 \pm 7.6$                        | 0.199            |
| LV diastolic parameters   |   |  |   |                                       |                  |
| E wave (cm/s)   | $80.6 \pm 19.7$   | $86.2 \pm 21.1$  | $80.7 \pm 21.7$   | $76.5 \pm 17.1$                       | 0.313            |
| A wave (cm/s)   | $56.2 \pm 15.1$   | $52.9 \pm 15.5$  | $55.8 \pm 12.7$   | $58.5 \pm 15.7$                       | 0.520            |
| F/A ratio   | 1.4(1.1-1.8)  | 1.7 (1.5–1.9)  | 1.4 (1.1–1.9)   | 1.3 (1.0–1.5)                         | 0.002            |
| Err rule<br>F' medial (cm/s)  | 9.5 (7.5–11.8)  | 10.7 (8.9–12.4)  | 9.3 (6.9–11.9)  | 8.7 (7.1-10.8)                        | 0.028            |
| E' lateral (cm/s)   | 134+33  | $148 \pm 30$   | 139 + 31  | $12.2 \pm 3.2$                        | 0.967            |
| E lateral (eni/s)   | 59(50-70)   | 60(50-70)  | 55(49-70)   | 12.2 = 3.2<br>6.0 (5.1-7.0)           | 0.363            |
| E/E lateral fatto   | J.7 (J.0-7.0)   | 0.0 (3.0-7.0)  | J.J (4.7-7.0)   | 0.0 (3.1 - 7.0)                       | ALC right strium |
| longitudinal strain; RV, right ventricle;<br>"Independent <i>t</i> -test, <i>p</i> -value or Mann-W | SD, standard deviation; TAPSE<br>Vhitney test, <i>p</i> -value for non-no | ge; LA, ien annular, FASF, J<br>2, tricuspid annular plane<br>formally distributed variabl | systolic excursion.<br>es. *Statistical significance de | noted by <i>p</i> -values < 0.05      | ALS, fight athum |

reduction in myocardial deformation, and RALS. Furthermore, this study demonstrates that RALS tended to decrease with age and males tended to have higher values compared to females, although this finding did not reach statistical significance.

Aging is associated with the development of myocardial fibrosis. Fibrotic tissue is stiffer and less compliant, resulting in subsequent cardiac dysfunction.<sup>37,38</sup> We hypothesised that a combination of stiff RA and RV diastolic dysfunction associated with age-related myocardial fibrosis will have poor myocardial deformation and this may result in an age-related decrease in RALS. However, we did not perform biomarkers or CMR imaging to objectively assess for the presence of fibrosis in this study.

The trend towards gender differences with RALS, although not statistically relevant, could be multifactorial in this study. In addition to a difference in biology, a higher BMI in females compared to males may explain these findings. Male participants were also younger and had higher RAVI and likely more compliant RA compared to females, which translated into higher RALS. Furthermore, males had a lower heart rate compared to females [65.0 (59.0–76.5) vs 76.0 (67.0–82.5) beats/min, p = 0.002], which allowed for a prolonged filling time of the RA

| Table 5. Multiple linear regression analysis for           predictors of RA longitudinal strain |                       |                   |         |                         |                   |         |                                 |
|---|-----------------------|-------------------|---------|-------------------------|-------------------|---------|---------------------------------|
|   | Univariate regression |                   |         | Multivariate regression |                   |         |                                 |
| Covariates  | Coef-<br>ficient      | Standard<br>error | p-value | Coef-<br>ficient        | Standard<br>error | p-value | <i>Multiple</i> r,<br>(p-value) |
| Age (years)   |                       |                   |         | -0.06                   | 0.10              | 0.558   | 0.40 (0.097)                    |
| Male gender   | 3.17                  | 2.13              | 0.141   | -1.15                   | 2.80              | 0.683   |                                 |
| BMI   | -0.43                 | 0.14              | 0.003   | -0.72                   | 0.29              | 0.013   |                                 |
| BSA   | -6.61                 | 6.30              | 0.297   | 16.20                   | 10.28             | 0.118   |                                 |
| RV S'   | 0.81                  | 0.51              | 0.114   | 0.47                    | 0.56              | 0.410   |                                 |
| RV base   | 0.0004                | 0.21              | 0.998   | -0.10                   | 0.22              | 0.644   |                                 |
| RV E'   | 0.49                  | 0.33              | 0.149   | 0.98                    | 0.73              | 0.186   |                                 |
| RV A'   | -0.11                 | 0.36              | 0.764   | -0.65                   | 0.75              | 0.387   |                                 |
| RV E'/A' ratio  | 2.11                  | 2.02              | 0.298   | -5.77                   | 5.76              | 0.319   |                                 |
| RA volume/<br>BSA ratio   | -0.19                 | 0.18              | 0.307   | -0.20                   | 0.19              | 0.307   |                                 |
| BMI, body mass index; BSA, body surface area; RA, right atrial; RV, right ventricular.          |                       |                   |         |                         |                   |         |                                 |

chamber and therefore increased stretch of the RA wall, with the resultant higher RALS.

Padeletti *et al.* did not find differences in RALS with regard to gender and ageing.<sup>10</sup> This lack of association between RALS, gender and aging may be related to the limited capacity of the software in identifying all the segments of the RA due to the higher tricuspid annulus deformation compared to the mitral valve. Furthermore, this may be attributed to varying sample sizes and racial differences in the two studies.

With recent advances in technology, Nemes *et al.*<sup>39</sup> and Qu *et al.*<sup>40</sup> using 3D STE and CMR, respectively, demonstrated obvious gender differences in RA strain. In contrast to our findings, they noted RA strain to be higher in females and showed an age-related decline in both genders.<sup>39,40</sup> The differences in analysis software and technique used in these studies may explain the discordances in results. Further studies are warranted to confirm our findings in a larger African population and also to further assess RALS.

We have confirmed the utility of RALS as a marker of subclinical disease in this population, as we did not see changes in RAVI with age but noted a trend towards lower RALS with increasing age. Therefore, RALS may anticipate RA impairment in disease prior to changes in traditional parameters such as RA size and volume. This may assist in earlier diagnosis of disease and prompt treatment strategies at a subclinical stage of the disease.

#### **Study limitations**

A minority of subjects were older than 60 years, due to the lower life expectancy in the South African population (the average life expectancy of an adult in 2014 was estimated at 59.1 years for males and 63.1 years for females, according to Statistics South Africa).<sup>41</sup> RA strain measurement values vary with different vendors and software packages and this needs to be taken into consideration when defining normal values. Exercise capacity of the study subjects was not assessed to unmask subclinical diastolic dysfunction and symptoms.

Since this was a secondary data analysis, sample size was restricted to what was collected in the parent study. A larger sample size might have been able to detect finer differences. The majority of the patients were obese or overweight and this is a reflection of the current 'normal' South African population. As this was a retrospective sub-study analysis, we did not have access to blood tests to screen for diseases, as these were not performed as part of the main study.

### Conclusion

We have presented the first normative values for RALS in a sub-Saharan African population. The normative data on RA strain and volumes according to age will help in differentiating normal from abnormal RA function and thus help in cardiovascular disease risk stratification in this population. Furthermore, this study provides a platform for future larger studies on RALS.

### References

1. Austin C, Alassas K, Burger C, Safford R, Pagan R, Duello K, et al. Echocardiographic assessment of estimated right atrial pressure and size predicts mortality in pulmonary arterial hypertension. *Chest* 2015; **147**(1): 198–208.

- Bouchardy J, Marelli AJ, Martucci G, Bottega N, Therrien J. Mirror image atrial dilatation in adult patients with atrial fibrillation and congenital heart disease. Int J Cardiol 2013; 167(3): 816–820.
- Sallach JA, Tang WHW, Borowski AG, Tong W, Porter T, Martin MG, *et al*. Right atrial volume index in chronic systolic heart failure and prognosis. *J Am Coll Cardiol Cardiovasc Imaging* 2009; 2(5): 527–534.
- Tadic M. The right atrium, a forgotten cardiac chamber: An updated review of multimodality imaging. J Clin Ultrasound 2015; 43(6): 335–345.
- Bommer W, Weinert L, Neumann A, Neef J, Mason DT, DeMaria A. Determination of right atrial and right ventricular size by two-dimensional echocardiography. *Circulation* 1979; 60(1): 91–100.
- Maceira AM, Cosin-Sales J, Prasad SK, Pennell DJ. Characterization of left and right atrial function in healthy volunteers by cardiovascular magnetic resonance. *J Cardiovasc Magn Reson* 2016; 18(1): 1–16.
- Sakata K, Uesugi Y, Isaka A, Minamishima T, Matsushita K, Satoh T, et al. Evaluation of right atrial function using right atrial speckle tracking analysis in patients with pulmonary artery hypertension. J Echocardiogr 2016; 14(1): 30–38.
- khedr L, Elasfar A, Hekal S, ElGendy E, Abdulaal M, Elsokkary H, et al. Assessment of left and right atrial geometrical changes in patients with stable coronary artery disease: Left and right atrial strain and strain rate imaging study. Egypt Heart J 2018; 70(2): 101–106.
- Haghighi ZO, Naderi N, Amin A, Taghavi S, Sadeghi M, Moladoust H, *et al.* Quantitative assessment of right atrial function by strain and strain rate imaging in patients with heart failure. *Acta Cardiol* 2011; 66(6): 737–742.
- Padeletti M, Cameli M, Lisi M, Zacà V, Tsioulpas C, Bernazzali S, *et al.* Right atrial speckle tracking analysis as a novel noninvasive method for pulmonary hemodynamics assessment in patients with chronic systolic heart failure. *Echocardiography* 2011; 28(6): 658–664.
- Hasselberg NE, Kagiyama N, Soyama Y, Sugahara M, Goda A, Ryo-Koriyama K, *et al.* The prognostic value of right atrial strain imaging in patients with precapillary pulmonary hypertension. *J Am Soc Echocardiogr* 2021; 34(8): 851–861.
- 12. Rudski LG, Lai WW, Afilalo J, Hua L, Handschumacher MD, Chandrasekaran K, *et al.* Guidelines for the echocardiographic assessment of the right heart in adults: a report from the American Society of Echocardiography. Endorsed by the European Association of Echocardiography, a registered branch of the European Society of Cardiology. *J Am Soc Echocardiogr* 2010; 23(7): 685–713.
- Lancellotti P, Badano LP, Lang RM, Akhaladze N, Athanassopoulos GD, Barone D, *et al.* Normal Reference Ranges for Echocardiography: rationale, study design, and methodology (NORRE study). *Eur Heart J Cardiovasc Imaging* 2013; 14(4): 303–308.
- 14. Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, et al. Recommendations for cardiac chamber quantification by echocardiography in adults: An update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. Eur Heart J Cardiovasc Imaging 2015; 16(3): 233–271.
- Mitchell C, Rahko PS, Blauwet LA, Canaday B, Finstuen JA, Foster MC, *et al.* Guidelines for performing a comprehensive transthoracic echocardiographic examination in adults: recommendations from the American Society of Echocardiography. *J Am Soc Echocardiogr* 2019; 32(1): 1–64.
- Soulat-Dufour L, Addetia K, Miyoshi T, Citro R, Daimon M, Fajardo PG, *et al.* Normal values of right atrial size and function according to age, sex, and ethnicity: results of the World Alliance Societies of Echocardiography Study. *J Am Soc Echocardiogr* 2021; 34(3): 286–300.

- Aurigemma GP, Gottdiener JS, Arnold AM, Chinali M, Hill JC, Kitzman D. Left atrial volume and geometry in healthy aging the cardiovascular health study. *Circ Cardiovasc Imaging* 2009; 2(4): 282–289.
- Lakatta EG, Levy D. Arterial and cardiac aging: major shareholders in cardiovascular disease enterprises: Part II: The aging heart in health: links to heart disease. *Circulation* 2003; **107**(2): 346–354.
- Meel R, Khandheria BK, Peters F, Libhaber E, Nel S, Essop MR. Left atrial volume and strain parameters using echocardiography in a black population. *Eur Heart J Cardiovasc Imaging* 2017; 18(3): 350–355.
- Dandel M, Lehmkuhl H, Knosalla C, Suramelashvili N, Hetzer R. Strain and strain rate imaging by echocardiography – basic concepts and clinical applicability. *Curr Cardiol Rev* 2009; 5(2): 133–148.
- Rai ABS, Lima E, Munir F, Faisal Khan A, Waqas A, Bughio S, *et al.* Speckle tracking echocardiography of the right atrium: the neglected chamber. *Clin Cardiol* 2015; **38**(11): 692–697.
- Padeletti M, Cameli M, Lisi M, Malandrino A, Zacã V, Mondillo S. Reference values of right atrial longitudinal strain imaging by twodimensional speckle tracking. *Echocardiography* 2012; 29(2): 147–152.
- Wang Y, Gutman JM, Heilbron D, Wahr D, Schiller NB. Atrial volume in a normal adult population by two-dimensional echocardiography. *Chest* 1984; 86(4): 595–601.
- Kou S, Caballero L, Dulgheru R, Voilliot D, De Sousa C, Kacharava G, et al. Echocardiographic reference ranges for normal cardiac chamber size: Results from the NORRE study. *Eur Heart J Cardiovasc Imaging* 2014; 15(6): 680–690.
- Nel S, Nihoyannopoulos P, Libhaber E, Essop MR, Ferreira dos Santos C, Matioda H, *et al.* Echocardiographic indices of the left and right heart in a normal black African population. *J Am Soc Echocardiogr* 2020; 33(3): 358–367.
- Karki DB, Pant S, Yadava SK, Vaidya A, Neupane DK, Joshi S. Measurement of right atrial volume and diameters in healthy nepalese with normal echocardiogram. *Kathmandu Univ Med J* 2014; **12**(2): 110–112.
- Grünig E, Henn P, D'Andrea A, Claussen M, Ehlken N, Maier F, *et al.* Reference values for and determinants of right atrial area in healthy adults by 2-dimensional echocardiography. *Circ Cardiovasc Imaging* 2013; 6(1): 117–124.
- D'Ascenzi F, Cameli M, Padeletti M, Lisi M, Zacà V, Natali B, *et al.* Characterization of right atrial function and dimension in top-level athletes: A speckle tracking study. *Int J Cardiovasc Imaging* 2013; 29(1): 87–94.
- 29. D'Oronzio U, Senn O, Biaggi P, Gruner C, Jenni R, Tanner FC, et al.

Right heart assessment by echocardiography: Gender and body size matters. J Am Soc Echocardiogr 2012; 25(12): 1251–1258.

- Peluso D, Badano LP, Muraru D, Dal Bianco L, Cucchini U, Kocabay G, et al. Right atrial size and function assessed with three-dimensional and speckle-tracking echocardiography in 200 healthy volunteers. Eur Heart J Cardiovasc Imaging 2013; 14(11): 1106–1114.
- Kengne AP, Bentham J, Zhou B, Peer N, Matsha TE, Bixby H, *et al.* Trends in obesity and diabetes across Africa from 1980 to 2014: An analysis of pooled population-based studies. *Int J Epidemiol* 2017; **46**(5): 1421–1432.
- Micklesfield LK, Kagura J, Munthali R, Crowther NJ, Jaff N, Gradidge P, et al. Demographic, socio-economic and behavioural correlates of BMI in middle-aged black men and women from urban Johannesburg, South Africa. *Glob Health Action* 2018; 11(suppl 2).
- Chirinos JA, De Buyzere M, Chahwala J, Satija V, Segers P, De Bacquer D, *et al*. Effect of obesity on left atrial strain in middle age: the Asklepios study. *J Am Coll Cardiol* 2018; **71**(11): A1608.
- Debari MK, Abbott RD. Adipose tissue fibrosis: Mechanisms, models, and importance. Int J Mol Sci 2020; 21(17): 1–24.
- Sokmen A, Sokmen G, Acar G, Akcay A, Koroglu S, Koleoglu M, *et al.* O impacto da obesidade isolada na função do ventrículo direito em adultos jovens. *Arq Bras Cardiol* 2013; **101**(2): 160–168.
- Csige I, Ujvárosy D, Szabó Z, Lorincz I, Paragh G, Harangi M, *et al.* The impact of obesity on the cardiovascular system. *J Diabetes Res* 2018; 2018.
- Donekal S, Venkatesh BA, Liu YC, Liu CY, Yoneyama K, Wu CO, *et al.* Interstitial fibrosis, left ventricular remodeling, and myocardial mechanical behavior in a population-based multiethnic cohort: The multi-ethnic study of atherosclerosis (MESA) study. *Circ Cardiovasc Imaging* 2014; 7(2): 292–302.
- Horn MA, Trafford AW. Aging and the cardiac collagen matrix: Novel mediators of fibrotic remodelling. J Mol Cell Cardiol 2016; 93: 175–185.
- Nemes A, Kormányos Á, Domsik P, Kalapos A, Ambrus N, Lengyel C, *et al.* Normal reference values of right atrial strain parameters using three-dimensional speckle-tracking echocardiography (results from the MAGYAR-Healthy study). *Int J Cardiovasc Imaging* 2019; 35(11): 2009–2018.
- Qu Y-Y, Buckert D, Ma G-S, Rasche V. Quantitative assessment of left and right atrial strains using cardiovascular magnetic resonance based tissue tracking. *Front Cardiovasc Med.* 2021; 8(June): 1–11.
- Statistics South Africa. Mid-year population estimates. online. www. statssa.gov.za/publications/./P03022014.pdf. 2014;(July):1–18.